

ANNUAL REPORT ON FY97 ONR SPONSORED RESEARCH

OCEAN DYNAMICS

Robert Pinkel

Scripps Institution of Oceanography

La Jolla California 92093-0213

phone: (619) 534-2056, fax:(619) 534-7132 email:rpinkel@ucsd.edu

Award #N00014-94-1-0046

LONG-TERM GOAL

To gain a more complete understanding of ocean dynamical processes, particularly at fine-scale, through intercomparison of high, mid- and low-latitude observations, both near the sea surface, in the main thermocline, and near the sea floor.

OBJECTIVES

To compare fine and internal wave scale measurements from equatorial, mid-latitude and Arctic observations. To quantify the relationship between fine-scale background conditions and the occurrence of microscale breaking.

APPROACH

Progress is effected through a steady-state cycle of instrument development, field observation and data analysis. The primary instruments employed include Doppler sonar and profiling CTD's. Generically, our instruments produce information which is quasi-continuous in space and time. Measurements typically span two decades in the wavenumber domain. This broad band space-time coverage enables the investigation of interactions which span space and time scales.

WORK COMPLETED

A multi-year study of shear, strain and Richardson number statistics has culminated in two publications in Journal of Physical Oceanography. The purely statistical approach was employed because, at fine-scale, many physically important quantities, such as the Vaisala frequency (inverse strain) are significantly non-Gaussian. A power spectrum provides an incomplete description of the field.

A focus of the effort was on a general model for ocean mixing. Bretherton, 1969, and Garrett and Munk, 1972, laid the groundwork for statistical mixing models. However, both efforts produced results which were extremely sensitive to the assumed levels of rms shear. Recent observational work by Gregg (1989) has suggested a more moderate dependence of

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Ocean Dynamics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, San Diego, Scripps Institution of Oceanography, La Jolla, CA, 92093				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

mixing on shear variance ($\epsilon \sim \langle s^2 \rangle^2$). Trading realism for simplicity, the GM type mixing formalism was revisited using correct pdf's of Richardson number and a more modern version of the internal wave spectrum (similar to Munk, 1981). With these advances, it proved straightforward to reconcile the logically pristine mixing formalism of GM 72 with the more recent observations.

RESULTS

Statistical predictions of mixing rates have been made in advance of an understanding of the detailed mechanisms involved. Numerous theoretical models, including the notable work of Henyey et al. (1986), have been posited, in the absence of quality observations of the sequence of events which lead to mixing. As a result of the statistical work, we are motivated to merge shear and density observations to obtain a phenomenological picture of ocean mixing. Mixing "events" are identified as density inversions of vertical scale 2 m and greater, as determined by a profiling CTD.

The study, based on 12 days of MBL data, resulted in several surprises. While there is some correspondence between mixing and the occurrence of low Richardson number, the strongest correspondence is with the occurrence of large vertical strain, $\frac{\partial \eta}{\partial z}$, and large rate of change of strain, $\frac{\partial w}{\partial z}$. These signals are associated with the propagation of small scale highly non-linear waves, which can be observed if the density field is examined in detail.

IMPACT/APPLICATION

We now know that small scale internal wave dynamics is intimately involved in determining when and where overturning occurs in the sea, and that Kelvin-Helmholtz instability is not the primary mixing mechanism. This knowledge can be applied in the analysis of existing data and in the prediction of other potential ocean mixing sites.

TRANSITIONS

The experimental capabilities developed in the course of this work are being applied in proposed investigations of coastal mixing, under ONR and NSF sponsorship. Sector-scan Doppler sonars, developed as an aspect of this work have been transitional for use in the near-shore experiments taking place at DUCK, N.C. Dr. Jerome Smith has lead in this effort.

RELATED PROJECTS

The Marine Boundary Layer observations benefit greatly from the technical legacy of preceding upper ocean programs such as SWAPP, LEADDEX and TOGA COARE. In these experiments the

instruments and techniques were developed which made MBL possible. Also, the Naval Sea Systems Command provided for a much needed overhaul of the Research Platform FLIP. The renovated FLIP performed quite well in support of MBL.

In turn, the instrument technology developed for MBL will be used in coming ONR research programs. SANDY DUCK, SHEBA and other programs will make use of the MBL systems. It is anticipated that the aspects of MBL which are unique geographically, such as the tidal and mixing observations of MBL I, can be used to plan a coming series of ONR experiments which hope to quantify the difference between the coastal ocean and the deep sea, at small scales.

REFERENCES

- Bretherton, F.P., 1969: Waves and turbulence in stably stratified fluids. *Radio Sci.*, 4, 1279-1287.
- Garrett, C.J.R., and W.H. Munk, 1972a: Space-time scales of internal waves. *Geophys. Fluid Dyn.*, 3, 225-264.
- Garrett, C.J.R., and W.H. Munk, 1972b: Ocean mixing by breaking internal waves. *Deep-Sea Res.*, 19, 823-832.
- Garrett, C.J.R., and W.H. Munk, 1975: Space-time scales of internal waves. A progress report. *J. Geophys. Res.*, 80, 291-297.
- Gregg, M.C., 1989: Scaling turbulent dissipation in the thermocline. *J. Geophys. Res.*, 94, 9686-9698.
- Pinkel, R., and S. Anderson, 1997: Shear, strain, and Richardson number variations in the thermocline. Part I: Statistical description. *J. Phys. Oceanogr.*, 27, (2) 264-281.
- Pinkel, R., and S. Anderson, 1997: Shear, strain, and Richardson number variations in the thermocline. Part II: Modeling Mixing. *J. Phys. Oceanogr.*, 27, (2) 282-290.